

EPCAM gene

Associated Syndrome Name: Lynch syndrome/Hereditary non-polyposis colorectal cancer (HNPCC)

EPCAM Summary Cancer Risk Table

CANCER	GENETIC CANCER RISK
Colorectal	High Risk
Endometrial	High Risk
Gastric	High Risk
Other	High Risk
Ovarian	High Risk
Skin	High Risk
Other	Elevated Risk
Pancreatic	Elevated Risk
Prostate	Elevated Risk

EPCAM gene Overview

Lynch syndrome ^{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11}

- Individuals with mutations in *EPCAM* have Lynch syndrome. This condition is also known as hereditary non-polyposis colorectal cancer (HNPCC).
- Men and women with Lynch syndrome due to mutations in *EPCAM* have a high risk of developing colorectal cancer, often at young ages. Colorectal cancer in patients with Lynch syndrome develops from adenomatous polyps which progress to cancer more quickly than polyps in individuals who do not have Lynch syndrome.
- Women with Lynch syndrome due to mutations in *EPCAM* have a high risk for developing endometrial and ovarian cancer, often at young ages.
- Patients with Lynch syndrome due to mutations in *EPCAM* are also believed to have an increased risk of developing a wide variety of other cancers, including small bowel, urinary tract, hepatobiliary tract, brain (usually glioblastoma), sebaceous gland, prostate, and pancreatic. These risks may be more significant in patients with a family history of these cancers. Therefore, the general screening and management recommendations provided below should be modified based on individualized risk assessment and counseling.
- Patients with Lynch syndrome due to mutations in *EPCAM* have an increased risk for gastric cancer. Earlier screening or more frequent intervals may be considered based on family history of upper gastrointestinal cancers or high-risk endoscopic findings. Random biopsy of the proximal and distal stomach should at minimum be performed on the initial upper endoscopy procedure to assess for *Helicobacter pylori*, autoimmune gastritis, and intestinal metaplasia.
- Studies have investigated the possibility that patients with Lynch syndrome have an increased risk for other cancers, including breast cancer and adrenocortical carcinoma. However, the data are not conclusive at this time and there are currently no medical management guidelines related to these cancers.
- Patients with Lynch syndrome have a high risk for developing second primary cancers following an initial diagnosis of colorectal or endometrial cancer. This includes a high risk for endometrial cancer in women following colorectal cancer and vice versa, a high risk for a second primary colorectal cancer in any portions of the colon or rectum remaining after surgical treatment, and a high risk for other Lynch associated cancers, such as those of the upper gastrointestinal tract, urinary tract, and other sites.
- Cancer risks for patients with Lynch syndrome due to mutations in *EPCAM* are currently estimated to be similar to those for patients with Lynch syndrome due to mutations in *MSH2*, and medical management guidelines are currently the same for patients with mutations in either gene. However, it is possible that this will change over time as we learn more about the exact risks associated with mutations in *EPCAM*. There are some early indications that endometrial cancer risk may be much lower for women with certain types of mutations in *EPCAM*.
- The timing of risk-reducing gynecological surgeries in individuals with Lynch syndrome due to mutations in *EPCAM* should be individualized based on whether childbearing is complete, the individual's medical and surgical history, family history, and other relevant factors.
- Although there are high risks for cancer in patients with Lynch syndrome, many of these risks can be greatly reduced with appropriate medical management. Guidelines for the medical management of patients with Lynch syndrome have been

developed by the National Comprehensive Cancer Network (NCCN) and other expert groups. These are listed below. It is recommended that patients with an *EPCAM* mutation and a diagnosis of Lynch syndrome be managed by a multidisciplinary team with expertise in medical genetics and the care of patients with this condition.

EPCAM gene Cancer Risk Table

CANCER TYPE	AGE RANGE	CANCER RISK	RISK FOR GENERAL POPULATION
Colorectal	To age 70 ^{1, 2, 12, 13, 14, 15}	33%-51%	1.9%
Endometrial	To age 70 ^{1, 2, 12, 13, 14, 15, 16}	21%-53%	1.9%
Ovarian	To age 70 ^{1, 2, 12, 13, 14, 15, 16}	8%-24%	0.6%
Urothelial	To age 70 ^{1, 2, 10, 14, 15, 17}	7%-21%	0.6%
Gastric	To age 70 ^{13, 14, 15, 16}	2%-9%	0.4%
Small Bowel	To age 70 ^{13, 14, 15, 16}	1%-3%	0.1%
Lynch-associated Skin Tumors	To age 70 ^{3, 8}	1%-9%	<1.0%
Overall cancer risk (Lynch cancers)	Risk for a second Lynch-related cancer after a first cancer diagnosis ^{18, 19}	Increased risk	NA
Prostate	To age 70 ^{2, 14, 15, 16}	13%-17%	6.3%
Brain	To age 70 ^{2, 13, 14, 15, 16}	2%-6%	0.3%
Pancreatic	To age 70 ^{11, 14, 15}	Up to 2%	0.6%
Hepatobiliary Tract	To age 70 ^{12, 13, 14, 15, 20}	Elevated risk	0.5%

EPCAM Cancer Risk Management Table

The overview of medical management options provided is a summary of professional society guidelines. The most recent version of each guideline should be consulted for more detailed and up-to-date information before developing a treatment plan for a particular patient.

This overview is provided for informational purposes only and does not constitute a recommendation. While the medical society guidelines summarized herein provide important and useful information, medical management decisions for any particular patient should be made in consultation between that patient and his or her healthcare provider and may differ from society guidelines based on a complete understanding of the patient's personal medical history, surgeries and other treatments.

CANCER TYPE	PROCEDURE	AGE TO BEGIN	FREQUENCY (UNLESS OTHERWISE INDICATED BY FINDINGS)
Colorectal	Colonoscopy ³	20 to 25 years, or 2 to 5 years younger than the earliest colorectal cancer diagnosis in family if it is under age 25	Every 1 to 2 years
	Consider the use of aspirin as a risk-reduction agent. ³	Individualized	Individualized
Endometrial	Patient education about the importance of quickly seeking attention for endometrial cancer symptoms, such as abnormal bleeding or menstrual cycle irregularities ³	Individualized	Individualized
	Consider endometrial biopsy. ³	30 to 35 years	Every 1 to 2 years
	Consider transvaginal ultrasound. ³	After menopause	Individualized

CANCER TYPE	PROCEDURE	AGE TO BEGIN	FREQUENCY (UNLESS OTHERWISE INDICATED BY FINDINGS)
	Consider hysterectomy. ³	40 to 50 years	NA
	Consider options for endometrial cancer risk-reduction agents (i.e. oral contraceptives, progestin intrauterine systems). ³	Individualized	NA
Ovarian	Consider bilateral salpingo-oophorectomy (BSO). ³	40 to 50 years	NA
	Consider options for ovarian cancer risk-reduction agents (i.e. oral contraceptives, progestin intrauterine systems). ^{3, 21}	Individualized	NA
Urothelial	Consider urinalysis. ³	30 to 35 years	Annually
Gastric	Test and treat for <i>Helicobacter pylori</i> infection. ³	Individualized	NA
	Upper endoscopy and consider extended duodenal exam, preferably performed in conjunction with colonoscopy. See clinical overview. ³	30 to 40 years, or earlier if there is a family history of gastric cancer at a young age	Every 2 to 4 years
Small Bowel	Upper endoscopy and consider extended duodenal exam, preferably performed in conjunction with colonoscopy. Push enteroscopy can be considered in place of upper endoscopy to enhance small bowel visualization. ³	30 to 40 years, or earlier if there is a family history of small bowel cancer at a young age	Every 2 to 4 years
Lynch-associated Skin Tumors	Consider skin exams. ³	Individualized	Every 1 to 2 years
Prostate	Consider prostate cancer screening with digital rectal examination (DRE) and prostate specific antigen (PSA). ^{3, 22}	40 years	Every 1 to 2 years, or adjusted based on PSA
Brain	Patient education about the importance of quickly seeking attention for signs and symptoms of neurologic cancer ³	Individualized	NA
Pancreatic	For patients with a family history of pancreatic cancer, consider available options for pancreatic cancer screening, including the possibility of endoscopic ultrasonography (EUS) and MRI/magnetic resonance cholangiopancreatography (MRCP). It is recommended that patients who are candidates for pancreatic cancer screening be managed by a multidisciplinary team with experience in screening for pancreatic cancer. ^{3, 23}	45 to 50 years, or 10 years younger than the earliest diagnosis of pancreatic cancer in the family	Annually
	Provide education about ways to reduce pancreatic cancer risk, such as not smoking and losing weight. ²⁴	Individualized	Individualized
Hepatobiliary Tract	Currently there are no specific medical management guidelines for hepatobiliary cancer risk in mutation carriers. ³	NA	NA
For Patients With A Cancer Diagnosis	For patients with a gene mutation and a diagnosis of cancer, targeted therapies may be available as a treatment option for certain tumor types (e.g., antibodies to PD-1) ²⁵	NA	NA

Information for Family Members

The following information for Family Members will appear as part of the MMT for a patient found to have a mutation in the EPCAM gene.

This patient's relatives are at risk for carrying the same mutation(s) and associated cancer risks as this patient. Cancer risks for females and males who have this/these mutation(s) are provided below.

Family members should talk to a healthcare provider about genetic testing. Close relatives such as parents, children, brothers and sisters have the highest chance of having the same mutation(s) as this patient. Other more distant relatives such as cousins, aunts, uncles, and grandparents also have a chance of carrying the same mutation(s). Testing of at-risk relatives can identify those family

members with the same mutation(s) who may benefit from surveillance and early intervention.

Parents who are concerned about the possibility of passing on an *EPCAM* mutation to a future child may want to discuss options for prenatal testing and assisted reproduction techniques, such as pre-implantation genetic diagnosis (PGD).³

References

1. Mu00f8ller P, et al. Cancer incidence and survival in Lynch syndrome patients receiving colonoscopic and gynaecological surveillance: first report from the prospective Lynch syndrome database. *Gut*. 2017 66:464-472. PMID: 26657901.
2. Dominguez-Valentin M, et al. Cancer risks by gene, age, and gender in 6350 carriers of pathogenic mismatch repair variants: findings from the Prospective Lynch Syndrome Database. *Genet Med*. 2020 22:15-25. PMID: 31337882.
3. Gupta S, et al. NCCN Clinical Practice Guidelines in Oncology® Genetic/Familial High-Risk Assessment: Colorectal, Endometrial, and Gastric. V 1.2025. Jun 13. Available at <https://www.nccn.org>.
4. Ryan S, et al. Risk of prostate cancer in Lynch syndrome: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2014 23:437-49. PMID: 24425144.
5. Kempers MJ, et al. Risk of colorectal and endometrial cancers in *EPCAM* deletion-positive Lynch syndrome: a cohort study. *Lancet. Oncol*. 2011 12:49-55. PMID: 21145788.
6. Grindedal EM, et al. Germ-line mutations in mismatch repair genes associated with prostate cancer. *Cancer Epidemiol Biomarkers Prev*. 2009 18:2460-7. PMID: 19723918.
7. Raymond VM, et al. Elevated risk of prostate cancer among men with Lynch syndrome. *J Clin Oncol*. 2013 31:1713-8. doi: 10.1200/JCO.2012.44.1238. Epub 2013 Mar 25. PMID: 23530095.
8. Idos G, Valle L. Lynch Syndrome. 2021 Feb 4. In: Pagon RA, et al., editors. *GeneReviews®* [Internet]. PMID: 20301390.
9. Lin KM, et al. Colorectal and extracolonic cancer variations in *MLH1/MSH2* hereditary nonpolyposis colorectal cancer kindreds and the general population. *Dis Colon Rectum*. 1998 41:428-33. PMID: 9559626.
10. Joost P, et al. Urinary Tract Cancer in Lynch Syndrome; Increased Risk in Carriers of *MSH2* Mutations. *Urology*. 2015 86:1212-7. PMID: 26385421.
11. Bogdanski AM, et al. Pancreatic cancer risk and survival in patients with Lynch syndrome: a nationwide Dutch cohort study. *EClinicalMedicine*. 2026 Jan 12;91:103755. PMID: 41583364.
12. Bonadona V, et al. Cancer risks associated with germline mutations in *MLH1*, *MSH2*, and *MSH6* genes in Lynch syndrome. *JAMA*. 2011 305:2304-10. PMID: 21642682.
13. Barrow E, et al. Cumulative lifetime incidence of extracolonic cancers in Lynch syndrome: a report of 121 families with proven mutations. *Clin Genet*. 2009 75:141-9. PMID: 19215248.
14. Mu00f8ller P, et al. Cancer risk and survival in path_MMR carriers by gene and gender up to 75 years of age: a report from the Prospective Lynch Syndrome Database. *Gut*. 2018 67:1306-1316. PMID: 28754778.
15. SEER*Explorer: An interactive website for SEER cancer statistics [Internet]. Surveillance Research Program, National Cancer Institute. [Cited 2026 May 5]. Available from <https://seer.cancer.gov/explorer/>.
16. Vasen HF, et al. Hereditary cancer registries improve the care of patients with a genetic predisposition to cancer: contributions from the Dutch Lynch syndrome registry. *Fam Cancer*. 2016 15:429-35. PMID: 26973060.
17. Skeldon SC, et al. Patients with Lynch syndrome mismatch repair gene mutations are at higher risk for not only upper tract urothelial cancer but also bladder cancer. *Eur Urol*. 2013 63:379-85. PMID: 22883484.
18. Win AK, et al. Risks of primary extracolonic cancers following colorectal cancer in lynch syndrome. *J Natl Cancer Inst*. 2012 104:1363-72. PMID: 22933731.
19. Win AK, et al. Risks of colorectal and other cancers after endometrial cancer for women with Lynch syndrome. *J Natl Cancer Inst*. 2013 105:274-9. PMID: 23385444.
20. Dowty JG, et al. Cancer risks for *MLH1* and *MSH2* mutation carriers. *Hum Mutat*. 2013 34:490-7. PMID: 23255516.
21. Daly M et al. NCCN Clinical Practice Guidelines in Oncology®: Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate. V 3.2026. Feb 19. Available at <https://www.nccn.org>.

22. Sprenkle PC, et al. NCCN Clinical Practice Guidelines in Oncology®: Prostate Cancer Early Detection. V 2.2026. Feb 18. Available at <https://www.nccn.org>.
23. Goggins M, et al. Management of patients with increased risk for familial pancreatic cancer: updated recommendations from the International Cancer of the Pancreas Screening (CAPS) Consortium. Gut. 2020 69:7-17. PMID: 31672839.
24. Ko AH, et al. NCCN Clinical Practice Guidelines in Oncology®: Pancreatic Adenocarcinoma. V 2.2026. Apr 22. Available at <https://www.nccn.org>.
25. https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/125514s031lbl.pdf.

Last Updated on 09-Jun-2026